

This is the list of all publications led by Ashutosh V. Kotwal or containing his significant contributions, *excluding* those related to the W boson mass measurement.

References

- [1] A. V. Kotwal, H. Kemeny, Z. Yang and J. Fan, “A low-latency graph computer to identify metastable particles at the Large Hadron Collider for real-time analysis of potential dark matter signatures,” *Sci. Rep.* **14**, no.1, 10181 (2024) doi:10.1038/s41598-024-60319-9
- [2] G. Aad *et al.* [ATLAS], “Combination of Searches for Higgs Boson Pair Production in pp Collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector,” *Phys. Rev. Lett.* **133**, no.10, 101801 (2024) doi:10.1103/PhysRevLett.133.101801 [arXiv:2406.09971 [hep-ex]].
- [3] G. Aad *et al.* [ATLAS], “Differential cross-section measurements of the production of four charged leptons in association with two jets using the ATLAS detector,” *JHEP* **01**, 004 (2024) doi:10.1007/JHEP01(2024)004 [arXiv:2308.12324 [hep-ex]].
- [4] G. Aad *et al.* [ATLAS], “Search for top-philic heavy resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *Eur. Phys. J. C* **84**, no.2, 157 (2024) doi:10.1140/epjc/s10052-023-12318-9 [arXiv:2304.01678 [hep-ex]].
- [5] K. Agashe, S. Airen, R. Franceschini, D. Kim, A. V. Kotwal, L. Ricci and D. Sathyan, “A new purpose for the W -boson mass measurement: Searching for New Physics in lepton+MET,” *Phys. Lett. B* **855**, 138774 (2024) doi:10.1016/j.physletb.2024.138774 [arXiv:2310.13687 [hep-ph]].
- [6] G. Aad *et al.* [ATLAS], “Observation of electroweak production of two jets and a Z-boson pair,” *Nature Phys.* **19**, no.2, 237-253 (2023) doi:10.1038/s41567-022-01757-y [arXiv:2004.10612 [hep-ex]].
- [7] A. V. Kotwal, “Searching for metastable particles using graph computing,” *Sci. Rep.* **11**, no.1, 18543 (2021) doi:10.1038/s41598-021-97848-6
- [8] S. V. Chekanov, A. V. Kotwal, C. H. Yeh and S. S. Yu, “Physics potential of timing layers in future collider detectors,” *JINST* **15**, no.09, P09021 (2020) doi:10.1088/1748-0221/15/09/P09021 [arXiv:2005.05221 [physics.ins-det]].
- [9] A. V. Kotwal, “A fast method for particle tracking and triggering using small-radius silicon detectors,” *Nucl. Instrum. Meth. A* **957**, 163427 (2020) doi:10.1016/j.nima.2020.163427 [arXiv:1910.14149 [physics.ins-det]].
- [10] C. H. Yeh, S. V. Chekanov, A. V. Kotwal, J. Proudfoot, S. Sen, N. V. Tran and S. S. Yu, “Studies of granularity of a hadronic calorimeter for tens-of-TeV jets at a 100 TeV pp collider,” *JINST* **14**, no.05, P05008 (2019) doi:10.1088/1748-0221/14/05/P05008 [arXiv:1901.11146 [physics.ins-det]].
- [11] M. Aaboud *et al.* [ATLAS], “Measurements of gluon-gluon fusion and vector-boson fusion Higgs boson production cross-sections in the $H \rightarrow WW^* \rightarrow e\nu\mu\nu$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector,” *Phys. Lett. B* **789**, 508-529 (2019) doi:10.1016/j.physletb.2018.11.064 [arXiv:1808.09054 [hep-ex]].

- [12] G. Aad *et al.* [ATLAS], “Search for high-mass dilepton resonances using 139 fb⁻¹ of pp collision data collected at $\sqrt{s}=13$ TeV with the ATLAS detector,” Phys. Lett. B **796**, 68-87 (2019) doi:10.1016/j.physletb.2019.07.016 [arXiv:1903.06248 [hep-ex]].
- [13] S. V. Chekanov, M. Beydler, A. V. Kotwal, L. Gray, S. Sen, N. V. Tran, S. S. Yu and J. Zuzelski, “Initial performance studies of a general-purpose detector for multi-TeV physics at a 100 TeV pp collider,” JINST **12**, no.06, P06009 (2017) doi:10.1088/1748-0221/12/06/P06009 [arXiv:1612.07291 [hep-ex]].
- [14] A. V. Kotwal, M. J. Ramsey-Musolf, J. M. No and P. Winslow, “Singlet-catalyzed electroweak phase transitions in the 100 TeV frontier,” Phys. Rev. D **94**, no.3, 035022 (2016) doi:10.1103/PhysRevD.94.035022 [arXiv:1605.06123 [hep-ph]].
- [15] B. Auerbach, S. Chekanov, J. Love, J. Proudfoot and A. V. Kotwal, “Sensitivity to new high-mass states decaying to $t\bar{t}$ at a 100 TeV collider,” Phys. Rev. D **91**, no.3, 034014 (2015) doi:10.1103/PhysRevD.91.034014 [arXiv:1412.5951 [hep-ph]].
- [16] A. V. Kotwal, H. Schellman and J. Sekaric, “Review of Physics Results from the Tevatron: Electroweak Physics,” Int. J. Mod. Phys. A **30**, no.06, 1541004 (2015) doi:10.1142/S0217751X15410043 [arXiv:1409.5163 [hep-ex]].
- [17] D. Bandurin, G. Bernardi, C. Gerber, T. Junk, A. Juste, A. Kotwal, J. Lewis, C. Mesropian, H. Schellman and J. Sekaric, *et al.* “Review of Physics Results from the Tevatron,” Int. J. Mod. Phys. A **30**, no.06, 1541001 (2015) doi:10.1142/S0217751X15410018 [arXiv:1409.4861 [hep-ex]].
- [18] C. Degrande, J. L. Holzbauer, S. C. Hsu, A. V. Kotwal, S. Li, M. Marx, O. Mattelaer, J. Metcalfe, M. A. Pleier and C. Pollard, *et al.* “Studies of Vector Boson Scattering And Triboson Production with DELPHES Parametrized Fast Simulation for Snowmass 2013,” [arXiv:1309.7452 [physics.comp-ph]].
- [19] J. Anderson, A. Avetisyan, R. Brock, S. Chekanov, T. Cohen, N. Dhingra, J. Dolen, J. Hirschauer, K. Howe and A. Kotwal, *et al.* “Snowmass Energy Frontier Simulations,” [arXiv:1309.1057 [hep-ex]].
- [20] G. Aad *et al.* [ATLAS], “Search for high-mass dilepton resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector,” Phys. Rev. D **90**, no.5, 052005 (2014) doi:10.1103/PhysRevD.90.052005 [arXiv:1405.4123 [hep-ex]].
- [21] G. Aad *et al.* [ATLAS], “Search for $t\bar{t}$ resonances in the lepton plus jets final state with ATLAS using 4.7 fb⁻¹ of pp collisions at $\sqrt{s} = 7$ TeV,” Phys. Rev. D **88**, no.1, 012004 (2013) doi:10.1103/PhysRevD.88.012004 [arXiv:1305.2756 [hep-ex]].
- [22] G. Aad *et al.* [ATLAS], “Search for resonances decaying into top-quark pairs using fully hadronic decays in pp collisions with ATLAS at $\sqrt{s} = 7$ TeV,” JHEP **01**, 116 (2013) doi:10.1007/JHEP01(2013)116 [arXiv:1211.2202 [hep-ex]].
- [23] G. Aad *et al.* [ATLAS], “Search for high-mass resonances decaying to dilepton final states in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector,” JHEP **11**, 138 (2012) doi:10.1007/JHEP11(2012)138 [arXiv:1209.2535 [hep-ex]].

- [24] G. Aad *et al.* [ATLAS], “A search for $t\bar{t}$ resonances in lepton+jets events with highly boosted top quarks collected in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector,” JHEP **09**, 041 (2012) doi:10.1007/JHEP09(2012)041 [arXiv:1207.2409 [hep-ex]].
- [25] G. Aad *et al.* [ATLAS], “A search for $t\bar{t}$ resonances with the ATLAS detector in 2.05 fb $^{-1}$ of proton-proton collisions at $\sqrt{s} = 7$ TeV,” Eur. Phys. J. C **72**, 2083 (2012) doi:10.1140/epjc/s10052-012-2083-1 [arXiv:1205.5371 [hep-ex]].
- [26] G. Aad *et al.* [ATLAS], “Search for dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector,” Phys. Rev. Lett. **107**, 272002 (2011) doi:10.1103/PhysRevLett.107.272002 [arXiv:1108.1582 [hep-ex]].
- [27] G. Aad *et al.* [ATLAS], “Search for high mass dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment,” Phys. Lett. B **700**, 163-180 (2011) doi:10.1016/j.physletb.2011.04.044 [arXiv:1103.6218 [hep-ex]].
- [28] T. Aaltonen *et al.* [CDF], “Search for High Mass Resonances Decaying to Muon Pairs in $\sqrt{s} = 1.96$ TeV $p\bar{p}$ Collisions,” Phys. Rev. Lett. **106**, 121801 (2011) doi:10.1103/PhysRevLett.106.121801 [arXiv:1101.4578 [hep-ex]].
- [29] T. Aaltonen *et al.* [CDF], “Improved Search for a Higgs Boson Produced in Association with $Z \rightarrow \ell^+\ell^-$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV,” Phys. Rev. Lett. **105**, 251802 (2010) doi:10.1103/PhysRevLett.105.251802 [arXiv:1009.3047 [hep-ex]].
- [30] T. Aaltonen *et al.* [CDF], “Measurement of the top quark mass with dilepton events selected using neuroevolution at CDF,” Phys. Rev. Lett. **102**, 152001 (2009) doi:10.1103/PhysRevLett.102.152001 [arXiv:0807.4652 [hep-ex]].
- [31] C. P. Hays, A. V. Kotwal and O. Stelzer-Chilton, “New Techniques in the Search for Z-prime Bosons and Other Neutral Resonances,” Mod. Phys. Lett. A **24**, 2387-2403 (2009) doi:10.1142/S021773230903179X [arXiv:0910.1770 [hep-ex]].
- [32] T. Aaltonen *et al.* [CDF], “A Search for high-mass resonances decaying to dimuons at CDF,” Phys. Rev. Lett. **102**, 091805 (2009) doi:10.1103/PhysRevLett.102.091805 [arXiv:0811.0053 [hep-ex]].
- [33] T. Aaltonen *et al.* [CDF], “A Search for the Higgs Boson Produced in Association with $Z \rightarrow \ell^+\ell^-$ Using the Matrix Element Method at CDF II,” Phys. Rev. D **80**, 071101 (2009) doi:10.1103/PhysRevD.80.071101 [arXiv:0908.3534 [hep-ex]].
- [34] T. Aaltonen *et al.* [CDF], “Search for the Higgs boson produced with $Z \rightarrow \ell^+\ell^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV,” Phys. Rev. Lett. **101**, 251803 (2008) doi:10.1103/PhysRevLett.101.251803 [arXiv:0807.4493 [hep-ex]].
- [35] A. Abulencia *et al.* [CDF], “Search for excited and exotic muons in the $\mu\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV,” Phys. Rev. Lett. **97**, 191802 (2006) doi:10.1103/PhysRevLett.97.191802 [arXiv:hep-ex/0606043 [hep-ex]].
- [36] D. Acosta *et al.* [CDF], “Search for excited and exotic electrons in the $e\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV,” Phys. Rev. Lett. **94**, 101802 (2005) doi:10.1103/PhysRevLett.94.101802 [arXiv:hep-ex/0410013 [hep-ex]].

- [37] D. Acosta *et al.* [CDF], “Search for long-lived doubly-charged Higgs bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV,” Phys. Rev. Lett. **95**, 071801 (2005) doi:10.1103/PhysRevLett.95.071801 [arXiv:hep-ex/0503004 [hep-ex]].
- [38] H. Gerberich, C. P. Hays and A. Kotwal, “Search for Excited and Exotic Muons at CDF,” [arXiv:hep-ex/0611006 [hep-ex]].
- [39] A. Kotwal [Tevatron Electroweak Working Group], “Status and plans of the Tevatron Electroweak Working Group,” AIP Conf. Proc. **753**, no.1, 149-157 (2005) doi:10.1063/1.1896697
- [40] C. P. Hays, Y. Huang, A. V. Kotwal, H. K. Gerberich, S. Menzemer, K. Rinnert, C. Lecci, M. Herndon and F. D. Snider, “Inside-out tracking at CDF,” Nucl. Instrum. Meth. A **538**, 249-254 (2005) doi:10.1016/j.nima.2004.08.104
- [41] D. Acosta *et al.* [CDF], “Search for doubly-charged Higgs bosons decaying to dileptons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV,” Phys. Rev. Lett. **93**, 221802 (2004) doi:10.1103/PhysRevLett.93.221802 [arXiv:hep-ex/0406073 [hep-ex]].
- [42] S. Hossenfelder, M. Bleicher, S. Hofmann, H. Stoecker and A. V. Kotwal, “Black hole relics in large extra dimensions,” Phys. Lett. B **566**, 233-239 (2003) doi:10.1016/S0370-2693(03)00835-9 [arXiv:hep-ph/0302247 [hep-ph]].
- [43] A. V. Kotwal and C. P. Hays, “Production and decay of spinning black holes at colliders and tests of black hole dynamics,” Phys. Rev. D **66**, 116005 (2002) doi:10.1103/PhysRevD.66.116005 [arXiv:hep-ph/0206055 [hep-ph]].
- [44] R. M. Thurman-Keup, A. V. Kotwal, M. Tecchio and A. Byon-Wagner, “W boson physics at hadron colliders,” Rev. Mod. Phys. **73**, 267-306 (2001) doi:10.1103/RevModPhys.73.267
- [45] U. Baur, R. Brock, J. Parsons, M. Albrow, D. Denisov, T. Han, A. Kotwal, F. Olness, J. Qian and S. Belyaev, *et al.* “Physics at Future Hadron Colliders,” eConf **C010630**, E4001 (2001) [arXiv:hep-ph/0201227 [hep-ph]].
- [46] A. V. Kotwal [CDF and D0], “Prospects for W boson physics in Run II at the Fermilab Tevatron,” Conf. Proc. C **000727**, 714-715 (2000)
- [47] A. V. Kotwal [D0 and CDF], “W boson physics at the Tevatron,” Conf. Proc. C **980723**, 467-471 (1998)
- [48] J. Yu and A. V. Kotwal, “Measurement of $\sin^2 \theta_W$ at the first muon collider,” AIP Conf. Proc. **435**, no.1, 398-405 (1998) doi:10.1063/1.56213
- [49] H. Schellman, A. Caldwell, L. de Barbaro, D. Harris, B. King, M. Klasen, A. Kotwal, D. Krakauer, S. Magill and K. S. McFarland, *et al.* “Deep inelastic scattering at a muon collider: Neutrino physics,” AIP Conf. Proc. **435**, no.1, 166-176 (1998) doi:10.1063/1.56230
- [50] M. R. Adams *et al.* [E665], “Proton and deuteron structure functions in muon scattering at 470-GeV,” Phys. Rev. D **54**, 3006-3056 (1996) doi:10.1103/PhysRevD.54.3006
- [51] C. M. Laymon, W. Amos, M. G. Burlein, H. T. Fortune, T. Ekenberg, A. Kotwal, J. M. O'Donnell, J. D. Silk, J. D. Zumbro and C. L. Morris, *et al.* “Pion elastic and inelastic $2(1)^+$ scattering on Ni-58, Ni-60, Ni-62, Ni-64 at $T(\pi) = 180$ -MeV,” Phys. Rev. C **53**, 1167-1172 (1996) doi:10.1103/PhysRevC.53.1167

- [52] M. Burlein, H. T. Fortune, W. M. Amos, T. L. Ekenberg, A. Kotwal, P. H. Kutt, J. M. O'Donnell, J. D. Silk, B. Boyer and A. Fuentes, *et al.* "PION INELASTIC SCATTERING TO THE LOW LYING BROAD 2+ IN NE-20," Phys. Rev. C **40**, 785-789 (1989) doi:10.1103/PhysRevC.40.785
- [53] J. M. O'Donnell, A. V. Kotwal and H. T. Fortune, "Three-state mixing among 0+ states in even Cd isotopes," Phys. Rev. C **38**, 2047-2051 (1988) doi:10.1103/PhysRevC.38.2047